arXiv:1305.0293v1 [nucl-th] 1 May 2013

Fission properties of the BCPM functional

Samuel A. Giuliani[*] and Luis M. Robledo[†]

Departamento de Fisica Teorica, Universidad Autonoma de Madrid, E-28049 Madrid, Spain

Abstract

We explore the properties of the Barcelona Catania Paris Madrid (BCPM) energy density functional concerning fission dynamics. Potential energy surfaces as well as collective inertias relevant in the fission process are computed for several nuclei where experimental data exists. Inner and outer barrier heights as well as fission isomer excitation energies are reproduced quite well in all the cases. The spontaneous fission half lives $t_{\mathrm{SF}}$are also computed using the standard semiclasical approach and the results are compared with the experimental data. A reasonable agreement with experiment is found over a range of 27 orders of magnitude but the theoretical predictions suffer from large uncertainties associated to the values of the parameters entering the spontaneous fission half life formula. The impact that increasing the pairing correlations strengths has in the spontaneous fission half lives is analyzed and found to be large in all the nuclei considered. Given the satisfactory description of the trend of fission properties with mass number we explore the fission properties of the even-even uranium isotope chain from ^{226}U to ^{282}U. Very large half lives are found when getting close to neutron number N=170.

---

[*] Electronic address: sam.and.giuliani@gmail.com

[†] Electronic address: luis.robledo@uam.es

I. INTRODUCTION

Fission is an unique physical phenomenon taking place in heavy atomic nuclei and involving the evolution of the system from its ground state to scission going through different intrinsic shapes that cover a wide range of different intrinsic deformation parameters [1–3]. Fission properties depend upon the competition between the surface energy term coming from the strong nuclear interaction and the Coulomb repulsion and therefore fission properties are often used as constraints to refine the parameters of effective nuclear interactions. Typical examples are the Gogny [4] D1S parametrization that was fine tuned to reproduce the fission barrier of ^{240}Pu [5]. More recently it has been used with Skyrme like interactions to define the UNEDEF1 parametrization [6, 7]. The gross features of fission can be understood from a mean field perspective involving the Hartree Fock Bogoliubov (HFB) theory [9, 30] and therefore it not surprising the large amount of studies devoted to this subject with Skyrme like interactions [7–10], Gogny ones [5, 11–17] or based on the relativistic mean field [18–20]. Fission observables also depend on the inertia of the system to the relevant collective degrees of freedom and therefore they are sensitive to pairing correlations. As a consequence, fission is a good testing ground to test both the theories and interactions commonly used in nuclear structure. In addition, the theoretical understanding of fission is relevant to other areas like energy production with nuclear reactors, radioactive waste degradation or the nucleosynthesis of heavy elements in the explosive galactic environments and following the r-process. Last but not least, a better understanding of fission could open the door to a better estimation of magic numbers and hence extra stability of super-heavy nuclei beyond Z=114. In this paper we explore the ability of a newly proposed energy density functional (EDF) denoted as Barcelona-Catania-Paris-Madrid (BCPM) [23] to describe fission. The BCPM [23] is a recent parametrization of the BCP EDF [21, 22, 24, 29] for nuclear structure calculations. Its free parameters have been adjusted to binding energies of even-even nuclei all over the nuclide chart, including deformed ones. Instead of the more traditional approaches, like Skyrme or Gogny like [9], where some central potential form is guessed (contact, gaussian, Yukawa, etc) and used afterwards to fit nuclear matter properties and/or the nuclear matter equations of state EoS (both symmetric and neutron), in the BCPM functional we start from a microscopic EoS that is fitted by means of a low order polynomial in the density. That polynomial fit is translated to a finite nuclei EDF just by replacing the nuclear matter density by the density of the finite nucleus. This procedure is inspired by the local density approximation LDA and is common practice in condense matter physics in practical applications of the Kohn-Sham theory. The EDF is complemented with a finite range surface term, a contact spin-orbit interaction of the same form as in popular Skyrme or Gogny forces, the Coulomb interaction and finally, a density dependent zero range pairing interaction [27] with strengths fitted to reproduce Gogny's neutron matter pairing gap. The parameters of the functional (essentially those of the finite range surface term plus some freedom in the polynomial fit to fine tune the binding energy per nucleon) are fitted to reproduce binding energies of the 518 even-even nuclei of the 2003 mass table evaluation of Audi and Wapstra. The properties of the interaction concerning quadrupole, octupole and fission dynamics have also been explored [23]. As shown below, the BCPM functional gives reasonable results for fission observables like spontaneous fission half lives, fission isomer excitation energies and inner and outer barrier heights. We have also shown that those results could be improved by slightly modifying the amount of pairing correlations, either by modifying the pairing strengths or by going beyond the mean field approximation to restore particle number symmetry broken by the HFB procedure. As a consequence of the satisfactory performance of BCPM in describing fission, we have explored fission properties of the uranium isotopic chain from proton drip line to the neutron drip line.

II. METHODS

The BCP energy density functional [24] is made of a bulk part which is inspired by fully microscopic and realistic calculations of symmetric and neutron matter equations of state [25, 26]. The two equations of state (symmetric and neutron matter) given as a a function of the nuclear density are parametrized using a fitting procedure in terms of low order polynomials of the densities. To account for finite size effects related to the surface energy, a phenomenological finite range interaction of gaussian type is included. In addition, the Coulomb interaction and the spin-orbit term are taken exactly as in the Skyrme or Gogny forces. To deal with open-shell nuclei we include in the BCP functionals a zero range density-dependent pairing interaction fitted to reproduce the nuclear matter gaps obtained with the Gogny force [27]. The calculations in finite nuclei are carried out with a modification of the HFBaxial computer code [28].

To describe fission we have resorted to the traditional procedure based on the mean field approach with pairing correlations: the Hartree- Fock- Bogoliubov (HFB) with constraints. As constraining operators we have used mainly the axially symmetric quadrupole moment operator $Q_{20} = z^2 - \frac{1}{2}(x^2 + y^2)$ although some exploratory calculations have also been performed with the octupole moment operator Q_{30} and the necking operator $Q_N(z_0) = \exp(-(z - z_0)^2/C_0^2)$. Axial symmetry has been preserved in the calculations because of the high computational cost involved in the release of this restriction. We are aware of the relevance of triaxiality specially in the height of the inner fission barrier but its effect is merely quantitative and to a much lesser extent, qualitative. On the other hand, reflection symmetry is allowed to break at any stage of the calculation leading to octupole deformed shapes and asymmetric fission. As a consequence of the breaking of the parity symmetry we are forced to constraint the center of mass to the origin as to prevent spurious solutions corresponding to a translation of the nucleus as a whole. The quasiparticle creation and annihilation operators of the HFB theory are expanded in a harmonic oscillator basis (HO) preserving axial symmetry and containing HO states with J_z quantum numbers up to 35/2 and up to 25 quanta in the z direction. The two lengths of the HO basis have been optimized in a few nuclei and the $b_\perp$ and b_z lengths obtained for each value of the quadrupole moment used in nearby nuclei. The minimization process uses the gradient method as described in [30, 31]. The most evident advantage of this method is its handling of constraints, which allows a larger number of them to be treated at once.

Dynamical parameters like quadrupole collective masses and moments of inertia are computed both in the so-called Adiabatic Time Dependent Mean Field Framework (ATDHFB) and the generator coordinate method (GCM) (see [35, 36] for specific details) . In both theories the evolution of the inertia parameters involve the inversion of huge matrices related to the stability matrix of the HFB theory and therefore we have recurred to the usual approximations of neglecting the off diagonal (corresponding to the interaction) terms in the stability matrix (cranking approximation [33, 34]). Zero point energy corrections to the HFB energy $\epsilon_0(Q_{20})$ are also considered in the ATDHFB and GCM approaches. In addition, the rotational energy correction computed following the phenomenological prescription of Ref [32] is also subtracted. This correction is very important to the shape of the potential energy as its value increases with deformation and can reach several MeV for large deformations.

The spontaneous fission half life is computed with the standard WKB formalism of quantum mechanics. In the WKB formalism the $t_{\rm sf}$ is given (in seconds) by the formula

$$t_{\rm sf} = 2.86\,10^{-21}(1+\exp(2S))$$

The action S along theQ_{20} constrained path is given by

$$S = \int_a^b dQ_{20}\sqrt{2B(Q_{20})(V(Q_{20}) - (E_{\rm GS} + E_0)}.$$

where the integration limits correspond to the classical turning points for the energy $E_{\rm GS} + E_0$. For the collective quadrupole inertia $B(Q_{20})$ we have used both the ATDHFB and the GCM expressions. The results obtained with the two different theories can differ in several orders of magnitude as the ATDHFB mass is known to be a factor in between 1.5 and 2 larger than the GCM mass. The potential energy $V(Q_{20})$ is given by the HFB mean field energy with two quantities subtracted: the zero point energy $\epsilon_0(Q_{20})$ associated to the quadrupole collective motion and the rotational energy correction given by the rotational approximation with the Yoccoz moment of inertia (see [32] for further details). Finally, an additional parameter E_0 is introduced in the action and is added to the ground state energy $E_{\rm GS}$. It is meant to represent the quantal ground state energy obtained after considering quantal fluctuations in the quadrupole degree of freedom around the ground state. This quantity could be estimated to be half of the square root of the curvature around the minimum divided by the collective inertia but it is often taken as a free parameter or kept fixed to some reasonable value. We have followed the later approach with $E_0 = 1.0$ MeV and estimated the impact of considering a larger value by repeating the calculations with $E_0 = 1.5$ MeV. The spontaneous fission half life obtained in this way is subject to several uncertainties that can lead to differences of several orders of magnitude. The uncertainties are: 1) the height of the inner fission barrier gets reduced when triaxiallity is considered in the mean field calculation. The amount of reduction is typically of a couple of MeV (but it can show some isotopic variation, see [11] for a recent account in the actinide region). 2) The value of E_0 can also make a difference in $t_{\rm SF}$ specially for long lived isotopes where the fission barrier is wide. The reason is that the value of E_0 alters the classical turning points. 3) The values of the correlation energy corrections to the HFB energy included in $V(Q_{20})$. The values are computed under certain assumptions and approximations and a better estimation of their values can lead to some changes to $V(Q_{20})$. 4) The approximations involved in the evaluation of the collective masses can lead to differences of the order of 40 or 50 % in their values 5) Finally, the amount of pairing correlations is also an important ingredient both in the evaluation of the zero point energy corrections as in the evaluation of the collective inertia. As shown below, changes of a few percent in the pairing strength values can lead to changes in the theoretical estimation of the half lives in the range of five to seven orders of magnitude.

On the other hand, the experimental values of the parameters defining the potential energy of the fission process, namely the inner and outer barrier heights and the excitation energy of the fission isomer seem to be more robust quantities to compare with as they are not as sensitive to pairing correlations as the other parameters. However, these experimental quantities are obtained under some model dependent assumptions that can mask the physical meaning of some of the parameters. Therefore, although we have compared our values with the experimental ones, we prefer to compare with spontaneous fission life times and more precisely with the evolution with Z and N of those quantities. We will also study the impact of changing various quantities entering the WKB formula.

III. RESULTS

A. Comparison with other interactions

Before comparing with the experimental data a comparison with the HFB results obtained with the two Gogny interactions, namely, D1S and D1M, is in order. The Gogny D1S interaction has been used in a thorough study of heavy nuclei properties, including fission, in Ref [11] and it has proved to reproduce quite nicely most of the properties analyzed. On the other hand, the fission properties of D1M have not been analyzed in detail but its good properties in many other aspects of nuclear structure like binding energies, radii and collective mode properties make it a good candidate to compare with. As we have already made comparison with D1S concerning fission properties [23] of actinides (^{240}Pu) and superheavies (^{262}Sg), we will just explore another actinide: the nucleus ^{234}U. In the lowest panel of Fig 1 we compare the HFB energy as a function of Q_{20} for the three functionals in the nucleus ^{234}U. We observe how the shape of the three curves look rather similar up $Q_{20} = 60$ b, apart from a constant shift of a few MeV, and from there one the D1M interaction produces a less pronounced decline than D1S and the BCPM results are in between but closer to D1M than



Figure 1: Comparison of HFB mean field quantities as a function of the mass quadrupole moment Q_{20} obtained with three different interactions. The BCPM EDF and the D1S and D1M parametrizations of the Gogny force. In the lowest panel, the HFB energy is given. In the middle panel the particle-particle correlation energy $E_{pp} = -\mathrm{Tr}(\Delta\kappa)$ is plotted for protons (dashed lines) and neutrons (full lines) for the three different kinds of calculations. In the panel above the octupole and hexadecapole deformation parameters are given. Finally, in the upper-most panel the ATDHFB collective inertia is given.

to D1S. In the next panel a quantity that can be used as a measure of pairing correlations, the particle-particle correlation energy given by $E_{pp} = -\mathrm{Tr}(\Delta\kappa)$, is depicted as a function of the mass quadrupole moment for both protons (dashed lines) and neutrons (full lines). Again the shape of the curves for protons and neutrons look rather similar in the three cases but there is again a shift in E_{pp} such that BCPM yields lower values than D1S and D1M. Those lower values correspond to less intense pairing correlations in the BCPM case being the quenching of proton pairing correlations more intense than the neutron ones. It is worth noticing that the pairing correlations in the proton channel for BCPM are significantly less intense that for the Gogny interactions. In the next panel, the octupole and hexadecapole moments as a function of Q_{20} are also given in the three cases. The results in the three cases are very similar, to the extent of appearing as a single curve for many Q_{20}values. Finally, in the upper-most panel the collective inertia in the ATDHFB approximation is plotted. As in the case of the HFB and particle-particle energies the shape of the curves are quite similar but the BCPM inertia is up to a factor of three larger than the inertias obtained with the Gogny forces. The large value of the BCPM inertia is a direct consequence of the quenched pairing correlations: less pairing correlations imply a lower gap and therefore smaller two quasiparticle energies. As the two quasiparticle energies enter the collective inertia in the denominator, quenched pairing correlations imply enlarged collective inertias. On the other hand, the D1M inertia is around 15% smaller than the D1S one, consistent again with the quenched pairing correlations in D1M as compared to the D1S ones. The GCM inertias, not depicted, look rather similar in shape to the ATDHFB ones but are a factor of 0.5-0.6 smaller. If the t_{SF} half life is computed with the WKB formula for the three cases with the GCM inertias, the values $t_{\mathrm{SF}} = 1.5 \times 10^{40}$s, 6.6×10^{34}s and 5.7×10^{23}s are obtained for BCPM, D1M and D1S, respectively. The large differences observed of up to 7 orders of magnitude can be attributed partly to the difference in the HFB curve but mostly to the different values of the collective inertias. The previous values have been obtained without taking into consideration the reduction of the inner barrier height as a consequence of triaxiallity. Also, increasing the value of the E_0parameter reduces the half lives by 6, 2 and 4 orders of magnitude respectively. Given the large variability of the half lives with the different parameters entering the WKB formula a direct comparison with the experimental data is meaningless and only comparisons along a series of nuclei, all of them computed with the same conditions, can lead to meaningful conclusions regarding fission properties.

B. Varying pairing strengths

In the BCPM functional the pairing interaction is taken as a density dependent contact pairing interaction with strength parameters fixed to reproduce the neutron matter pairing gap of the Gogny force [27]. We have shown in the previous subsection that the particle-particle correlation energy, a quantity related to the amount of pairing correlations, was much smaller for BCPM than for the Gogny forces leading to much larger collective inertias. It is therefore reasonable to investigate the behavior of fission properties as a function of the pairing strength. To this end, a parameter η has been introduced as a multiplication factor in front of the pairing gap field Δ_{kl}. For the sake of simplicity we have considered an unique parameter for both protons and neutrons although different parameters will give more freedom to adjust to experimental data. The outcome of the calculations with η values of 1.05 and 1.10 for the nucleus ^{234}U are presented in Fig 2. We observe in the lowest panel how increasing the pairing strength by 10 % ($\eta = 1.10$) we gain of the order of 1 MeV in binding energy and the gain is even larger on top of the inner barrier. The net effect is to slightly decrease the inner barrier eight (B_I) whereas the other parameters, namely the outer barrier height B_{II} and the fission isomer excitation energy remain more or less the same. The particle-particle correlation energies E_{pp} for protons and neutrons increase with increasing η but the slope is larger for neutrons than for protons. The multipole moment values do not change at all in they are on top of each other for different η values. Finally, the impact on the collective inertia is clearly visible. Increasing the pairing strength by 5 % reduces the collective inertia by roughly 30 % whereas a 10 % increase leads to a reduction of a 50 %. The impact on the t_{SF} is dramatic, decreasing by 11 orders of magnitude in going from $\eta = 1.0$ to $\eta = 1.05$ and six additional orders of magnitude in going from $\eta = 1.05$ to $\eta = 1.10$. This result is a clear indication of the very important role played by pairing correlations in the description of fission. The result suggests that experimental fission data could be used to fine tune the pairing strength instead of more traditional approaches based on odd-even staggerings. From a theoretical perspective the result also points to the very important role that the correlations associated to particle number symmetry restoration should have in fission dynamics. Restoring particle number symmetry usually leads to larger pairing correlations than the ones present at the mean field level and therefore will have a tremendous impact on fission half lives. In this respect it is worth mentioning that the dependence on density of BCPM is on integer powers of the density allowing the use of the regularization techniques suggested to solve some technical problems associated to the evaluation of energy kernel overlaps typical of symmetry restoration theories (see Refs [37, 38]and references therein).

C. Nuclei with known experimental data

In order to validate the BCPM functional as a functional able to describe fission, we have performed calculations for those even-even nuclei where the spontaneous fission half life has been measured. We will also compare the parameters defining the theoretical potential energy surface, namely the inner and our barrier heights (B_I and B_{II}) and the excitation energy of the fission isomer E_{II} with available experimental data. It has to be mentioned that the experimental data for B_I and B_{II} is model dependent and therefore less reliable that the pure t_{SF} data. Recently, a thorough comparison of theoretical results with the experimental data has been carried out in Ref. [7]. In table I the experimental and theoretical values for B_I, B_{II} and E_{II} are given for all nuclei where experimental data exists [39, 40]. The theoretical values have been obtained by considering the HFB energy as a function of Q_{20} with the rotational energy correction (computed in the way described in the previous section) subtracted. The effect of the zero point energy correction $\epsilon_0(Q_{20})$ has not been included mainly because it is almost constant as a function of Q_{20}. We notice that the theoretical predictions for B_I are typically one or two MeV larger than experiment. This is not surprising as it is well known that the inner fission barrier is affected by triaxiallity and its height typically decreases by one or two MeV when triaxiallity is included in the calculation. Triaxiallity is not included at present because we still do not have access to a triaxial code incorporating the BCPM functional but work in this direction is in progress. The situation is slightly better in the comparison with the E_{II} and B_{II}values. For them, no triaxial effects are expected and the agreement is better than for the B_I.

In table II the theoretical results obtained for different choices of the E_0 and η parameters are compared to the known experimental values. The experimental t_{SF} values span a range of 27 orders of magnitude for a mass range $A = 232 - 286$. The theoretical predictions, not including triaxial effects and computed with the GCM masses and zero point energies, span an even larger range of values and show a large variability depending upon the choices for the

Nucleus	B_I(th)	E_{II}(th)	B_{II}(th)	B_I(exp)	E_{II}(exp)	B_{II}(exp)
^{234}U	5.87	1.78	5.59	4.80	–	5.50
^{236}U	6.49	1.90	6.04	5.0	2.75	5.67
^{238}U	6.99	2.03	6.54	6.30	2.55	5.50
^{238}Pu	6.91	1.85	5.20	5.60	2.4	5.10
^{240}Pu	7.43	2.08	5.69	6.05	2.8	5.15
^{242}Pu	7.72	2.27	6.30	5.85	2.2	5.05
^{244}Pu	7.89	2.47	6.30	5.70	–	4.85
^{240}Cm	6.8	1.2	3.90	–	2	–
^{242}Cm	7.4	1.7	4.5	6.65	1.9	5.0
^{244}Cm	8.0	1.9	5.0	6.18	2.2	5.10
^{246}Cm	8.4	2.3	5.5	6.0	–	4.80
^{248}Cm	8.34	2.04	5.47	5.80	–	4.80
^{250}Cf	8.65	1.25	4.24	–	–	3.8
^{252}Cf				–	–	3.5

Table I: Fission barrier height parameters B_I (inner) and B_{II}(outer) as well as excitation energy of the fission isomer E_{II}. The three parameters are given in MeV. The theoretical values have been obtained from the rotational energy corrected HFB potential energy surface. The experimental values are taken from [39] for the E_{II} and from [40] for the B's.

parameters. Focusing on the "standard" theoretical values $\eta = 1.0$ and $E_0 = 1.0$ we observe differences of up to 16 orders of magnitude for the lighter nuclei that decrease to differences of just a couple of orders of magnitude for the heavier ones. The largest differences are observed for nuclei with higher and wider barriers where the impact of parameters like E_0 is larger. The comparison in isotopic chains indicate that the behavior with neutron number is not as bad as the absolute values. The same conclusion can be extracted from the overall trend with mass number obtained from the table. Therefore, we conclude that the HFB predictions although subject to large uncertainties due to uncontrolled approximations in the evaluation of the different parameters can be used to guess with a reasonable precision the trends of t_{SF} with mass number.

Spontaneous Fission Half-Life Times (s)						
		E_0= 1.0 MeV			E_0= 1.5 MeV	experimental
	Nuclei	η=1.10	η=1.05	η=1.00	η=1.0	arXiv-1302.7117
1	232U	3.842e+23	4.213e+29	2.275e+37	6.761e+30	2.5e+21
2	234U	7.211e+23	1.180e+30	9.987e+37	1.621e+31	4.7e+23
3	236U	8.697e+23	2.205e+29	9.759e+34	1.744e+28	7.8e+23
4	238U	4.018e+23	7.371e+28	7.117e+34	7.160e+28	2.6e+23
5	240Pu	3.107e+14	8.901e+19	3.567e+26	1.269e+21	1.5e+18
6	248Cm	8.684e+6	1.798e+11	3.707e+16	8.413e+12	1.3e+14
7	250Cf	2.747e+5	3.534e+10	1.074e+17	2.048e+13	5.2e+11
8	250Fm	1.719e-1	2.937e+3	5.098e+8	3.742e+4	2.6e+7
9	252Fm	1.006e00	3.971e+4	4.810e+9	3.189e+5	4.0e+9
10	254Fm	4.648e-5	1.941e-1	1.181e+3	5.037e-2	1.9e+7
11	256Fm	3.327e-6	1.932e-3	2.910e+1	2.447e-2	1.0e+4
12	252No	1.090e-6	9.505e-3	2.782e+2	8.740e-2	1.2e+1
13	254No	5.490e-5	6.689e-1	3.377e+4	1.118e+1	3.0e+4
14	256No	3.853e-5	1.097e-1	1.424e+2	2.071e-2	1.1e+2
15	256Rf	1.355e-5	1.050e+1	1.277e+8	3.337e+4	6.4e-3
16	258Rf	3.157e-5	2.879e00	9.416e+5	3.892e+2	1.4e-2
17	260Rf	5.623e-5	6.264e-1	6.856e+4	4.015e+1	5.1e-2
18	258Sg	1.042e-7	5.609e-5	9.134e-2	1.611e-3	5.2e-3
19	260Sg	8.999e-7	6.061e-4	8.862e-1	1.517e-2	7.2e-3
20	262Sg	9.885e-6	8.554e-3	2.169e+1	4.045e-1	7.0e-3
21	264Hs	3.585e-7	8.266e-4	2.735e+1	5.065e-1	1.6e-3
22	286Fl	2.502e-10	1.039e-5	2.625e00	1.724e-3	1.3e-1

Table II: Spontaneous fission half lives for a set of nuclei. Different theoretical predictions corresponding to different choices of E_0 and η are given. The experimental data is taken from [41] See text for discussion.

D. Neutron rich uranium isotopes

In the previous section we concluded that the description of fission based on the HFB theory is subject to large uncertainties coming from a poor understanding on the way the different quantities should be computed. However, we also concluded that the HFB theory is giving reasonable well the t_{SF} trends with mass number. Based on this conclusion, we have performed calculations in the uranium isotopic chain from the light uranium ^{226}U up to the neutron drip line corresponding to ^{282}U with the aim of understanding and analyzing the trends in spontaneous fission half lives. In Fig 3 the HFB potential energies are depicted as a function of Q_{20} for the isotopes of uranium ^{240}U, ^{250}U, ^{260}U, ^{270}U and ^{280}U. The energies of the heavier isotopes have been shifted by different amounts of energy (0, 55, 100, 135 and 160 MeV, respectively) to fit all the plots in a single plot.

We observe the ground state evolution from a quadrupole deformed ground state in ^{240}U with $\beta_2 = 0.26$ to an spherical one for ^{270}U (corresponding to $N = 178$) that is still spherical for ^{280}U. It is also worth mentioned the existence of a second fission isomer in ^{240}U and ^{250}U that disappears in ^{260}U, where the first fission isomer is shifted to larger quadrupole moment values and zero octupole moment. Two fission isomers reappear in ^{270}U but both are located at a very high excitation energy and different quadrupole deformations than the ones in the light uranium isotopes. As a consequence of the higher and wider barriers as the neutron number approaches the neutron drip line we expect increasing t_{SF} values as can be observed in table III. A decrease in the t_{SF} values is observed up to mass number $A = 256$ but from there on a steady increase with mass number is observed. The increase in t_{SF} is very large for ^{262}U that corresponds to a neutron number of 170. From this isotope up to the last one considered the values of t_{SF} are very large exceeding in more than 20 order of magnitude the typical values of the isotopes close to the stability line. As in previous cases, the t_{SF} values have been obtained with the GCM collective mass and not exploring the consequences of triaxiallity in the first barrier.

IV. CONCLUSIONS

The fission properties of several actinides and super-heavy nuclei have been computed with the EDF BCPM recently proposed. The theoretical results for the spontaneous fission

Uranium Spontaneous Fission Half-Life Times (s)

	E_0= 1.0 MeV			E_0= 1.5 MeV	Experimental
Nucleus	η=1.10	η=1.05	η=1.00	η=1.0	
226U			1.193e+26	1.516e+19	
228U			1.678e+33	1.622e+26	
230U			1.724e+35	3.739e+28	
232U	3.842e+23	4.213e+29	2.275e+37	6.761e+30	2.5e+21
234U	7.211e+23	1.180e+30	9.987e+37	1.621e+31	4.7e+23
236U	8.697e+23	2.205e+29	9.759e+34	1.744e+28	7.8e+23
238U	4.018e+23	7.371e+28	7.117e+34	7.160e+28	2.6e+23
240U			8.151e+34	2.059e+29	
242U			4.425e+35	1.109e+30	
244U			3.414e+32	4.121e+26	
246U			5.066e+28	3.068e+21	
248U			1.266e+26	3.943e+19	
250U		2.601e+23	2.776e+27	9.389e+19	
252U			4.862e+29	1.846e+21	
254U			1.796e+29	5.970e+21	
256U			5.948e+27	2.473e+20	
258U			1.786e+29	3.608e+22	
260U		1.058e+26	4.071e+31	2.577e+24	
262U			3.416e+43	1.952e+36	
264U			4.781e+49	4.567e+42	
266U			4.921e+58	7.239e+51	
268U			5.448e+68	7.013e+61	
270U		4.314e+69	3.123e+76	1.317e+68	
272U			1.126e+83	1.680e+75	
274U			7.609e+84	6.402e76	
276U			5.363e+98	7.268e+89	
278U			1.600e+86	3.652e+78	
280U		1.926e+69	4.209e+74	1.109e+67	
282U			3.872e+63	3.351e+56	

Table III: Spontaneous fission half lives for all the uranium isotopes from the proton ($A = 226$) to

half lives show a large variability consequence of uncertainties in the evaluation of some parameters of the theory and also on the strong dependence of the collective inertia with pairing correlations. As a consequence of the large uncertainties in the theoretical results we are only able to compare with the experimental data trend with mass number (for instance the reduction by 27 orders of magnitude in the spontaneous fission half lives in going from $A = 232$ to $A = 286$). The theoretical predictions seem to reproduce such trend giving us confidence in the relevance of the study of fission barrier properties of neutron rich uranium isotopes. There we find that the spontaneous fission half lives remain more or less constant up to A=260 where they increase enormously as a consequence of the proximity to the neutron number N=170 that is predicted to be a new magic number. On the other hand, a comparison of the parameters defining the potential energy surface for fission (inner and out barrier heights and fission isomer excitation energies) with the model dependent "experimental data" show a rather good agreement that gives us additional confidence on the validity of our conclusions. The results obtained clearly show that more attention has to be paid to a proper description (including beyond mean field effects) of pairing correlations in the configurations relevant to fission.

Acknowledgments

Work supported in part by MICINN grants Nos. FPA2009-08958, FIS2009-07277 and by the Consolider-Ingenio 2010 program CPAN CSD2007-00042 and MULTIDARK CSD2009-00064.

---

[1] H.J. Krappe and K. Pomorski, Theory of Nuclear Fission, Lect. Notes in Phys. **838** (2012).

[2] D.N. Poenaru, *Nuclear decay modes*, Institute of Physics Publishing (1996).

[3] Hans J. Specht, Rev. Mod. Phys. **46**, 773 (1974); S. Björnholm and J. E. Lynn, Rev. Mod. Phys. **52**, 725 (1980); "Fifty Years of Fission", Nucl. Phys. **A502** (1989).

[4] J. Decharge and D. Gogny, Phys. Rev. C**21**, 1568 (1980).

[5] J.F. Berger, M. Girod and D. Gogny. Nucl. Phys. **A428**, 23c (1984).

[6] N. Nikolov, N. Schunck, W. Nazarewicz, M. Bender and J. Pei, Phys Rev **C83** 034305 (2011).

[7] J. McDonnell, N. Schunck and W. Nazarewicz, arXiv: 1301.7587

[8] J. Erler, et al Phys. Rev **C85**, 025802 (2012).

[9] M. Bender, P.-H. Heenen, and P.-G. Reinhard, Rev. Mod. Phys. **75**, 121 (2003).

[10] A. Staszczak, A. Baran and W. Nazarewicz, arXiv: 1208.1215

[11] J.-P. Delaroche, M. Girod, H. Goutte and J. Libert, Nucl. Phys. **A 771, 103** (2006).

[12] N. Dubray, H. Goutte, and J.-P. Delaroche, Phys. Rev. C**77**, 014310 (2008).

[13] V. Martin and L.M. Robledo, Int. J. Mod. Phys. **E 18**, 788 (2009).

[14] S. Perez-Martin and L.M. Robledo, Int. J. Mod. Phys. **E 18**, 861 (2009).

[15] W. Younes and D. Gogny, Phys. Rev. **C 80**, 054313 (2009).

[16] J.L. Egido and L.M. Robledo, Phys. Rev. Lett. **85**, 1198 (2000).

[17] M. Warda, J.L. Egido, L.M. Robledo and K. Pomorski, Phys. Rev. C**66**, 014310 (2002); Intl. J. of Mod. Phys. E**13**, 169 (2004).

[18] H. Abusara, A.V. Afanasjev and P. Ring Phys. Rev. **C82**, 044303 (2010).

[19] B.-N. Lu, E.-G. Zhao and S.-G. Zhou, Phys. Rev. C85, 11301 (2012).

[20] A.V. Afanasjev, arXiv: 1303.1206

[21] L.M. Robledo, M. Baldo, P. Schuck and X. Viñas, Phys. Rev. **C77**, 051301(R) (208).

[22] L.M. Robledo, M. Baldo, P. Schuck and X. Viñas, Phys. Rev. **C81**, 034315 (2010).

[23] M. Baldo, L.M. Robledo, P. Schuck and X. Viñas, arXiv: 1210.1321 (2012).

[24] M. Baldo, P. Schuck, and X. Viñas, Phys. Lett. B **663**, 390 (2008).

[25] M. Baldo, C. Maieron, P. Schuck, and X. Viñas, Nucl. Phys. **A736**, 241 (2004).

[26] G. Taranto, M. Baldo and G.F. Burgio, arXiv: 1302.6882 (2013).

[27] E. Garrido, P. Sarriguren, E. Moya de Guerra, and P. Schuck, Phys. Rev. C **60**, 064312 (1999).

[28] L.M. Robledo, *HFBaxial computer code* (2002).

[29] M. Baldo, L.M. Robledo, P. Schuck, and X. Viñas J. of Phys. **G37** 064015 (2010)

[30] P. Ring and P. Shuck, *The Nuclear Many Body Problem* (Springer–Verlag Edt. Berlin, 1980).

[31] L.M. Robledo and G.F. Bertsch, Phys. Rev **C84**, 014312 (2011).

[32] J.L Egido and L.M. Robledo, Lecture Notes in Physics **641**, 269 (2004).

[33] M. Girod and B. Grammaticos, Nucl Phys. **A330**, 40 (1979)

[34] M.J. Giannoni and P. Quentin, Phys. Rev. **C21** (1980) 2060; *ibid* 2076

[35] A. Baran, et al., Phys Rev **C84**, 054321 (2011).

[36] J. Libert, M. Girod, and J.-P. Delaroche, Phys. Rev. **C60**, 054301 (1999).

[37] L.M. Robledo, Intl. J. of Mod. Phys. **E16** 337 (2007).

[38] L.M. Robledo, Jour. of Phys.**G37** 064020 (2010).

[39] B. Singh, R. Zywina, and R. Firestone, Nucl. Data Sheeets **97**, 241 (2002).

[40] R. Capote et al Nucl. Data Sheets **110**, 3107 (2009).

[41] N.E. Holden and D.C. Hoffman, Pure Appl. Chem, **72**, 1525 (2000).



Figure 2: Same as figure 1 but for different pairing strengths. The pairing strengths are given in terms of the reference value and a scaling parameter η taking the values 1.00 (the standard calculation), 1.05 and 1.10.



Figure 3: HFB energies for some neutron rich uranium isotopes as a function of the quadrupole moment.